



# **ORIGINAL:** Comparison of Knowledge and Attitude among Nurses in Special Wards and Emergency of the Hospital about COVID-19: A Cross-Sectional Study in the North of Iran

**Fereshteh Araghian** School of Nursing and Midwifery, Mazandaran University of Medical Sciences, Sari, Iran. Mojarad Hamid Salehinia Social Determinants of Health Research Center, Birjand University of Medical Sciences, Birjand, Iran. Mohammad Ali Heydari Department of Medical Surgical Nursing, School of Nursing and Midwifery, Mazandaran University of Gorji Medical Sciences, Sari, Iran. Abolfazl Hosseinnataj Department of Biostatistics and Epidemiology, School of Health, Mazandaran University of Medical Sciences, Sari, Iran. Fatemeh Talebian Student Research Committee, School of Nursing and Midwifery, Mazandaran University of Medical Sciences, Sari, Iran. Traditional and Complementary Medicine Research Center, Addiction Institute, Mazandaran University **Tahereh Yaghoubi** of Medical Sciences, Sari, Iran. Leila Jouybari Nursing Research Center, Golestan University of Medical Sciences, Gorgan, Iran.

## **ARTICLE INFO**

Submitted:	10 Jun 2021
Accepted:	14 Jul 2021
Published:	01 Sep 2021

Keywords: Attitude; COVID-19; Knowledge; Nurses

#### Correspondence:

TaherehYaghoubi,SchoolofNursingandMidwifery,MazandaranUniversityofMedicalSciences,Sari,Iran.Email:tyaghubi@mazums.ac.irORCID:0000-0001-8526-6745

#### **Citation:**

Araghian Mojarad F, Salehinia H, Heydari Gorji MA, Hosseinnataj A, Talebian F, Yaghoubi T, Jouybari L. Comparison of Knowledge and Attitude among Nurses in Special Wards and Emergency of the Hospital about COVID-19: A Cross-Sectional Study in the North of Iran. Tabari Biomed Stu Res J. 2021;3(3):14-20.

doi 10.18502/tbsrj.v3i3.6928

### ABSTRACT

**Introduction:** There is a lack of studies on the knowledge and attitudes of health care workers towards COVID-19. The aim of this study was to compare the knowledge and attitudes of nurses in emergency departments and special wards of medical training centers in Mazandaran province about COVID-19 in 2020.

**Material and Methods:** The present study is a descriptive crosssectional study. Which was performed in the emergency and special wards of adult educational and medical centers of Mazandaran province in 2020. Sampling is census that is easily accessible and non-random. Data collection tools included demographic information form and a researcher-made questionnaire related to knowledge and attitude. Data were analyzed by analytical tests using SPSS version 16.

**Results:** The findings of this study showed that the level of nurses 'knowledge was  $7.78 \pm 0.85$  and the level of nurses' attitude was  $6.87\pm 0.90$ . There was no statistically significant difference between knowledge and attitude in emergency and special ward's nurses. There is no significant relationship between education level, age and work experience with knowledge and attitude.

**Conclusion:** According to the findings, the level of knowledge of nurses was insufficient and the level of attitude of nurses was sufficient. Therefore, it is recommended nursing managers should plan to increase the level of awareness of nursing staff. Prepare virtual educational content in the field of treatment, nursing care and complications of COVID-19.

# Introduction

oronaviruses are RNA viruses that are widely distributed among humans,

other mammals, and birds and cause respiratory, intestinal, liver, and neurological

diseases (1, 2). Six species of coronavirus are known to be a human disease (3). Four viruses of - E229, OC43, NL63, and HKU1 are common and typically show cold symptoms in individuals with immune system defects (4). The other two strains -Acute Respiratory Coronavirus Syndrome (SARS-CoV) and Middle East Respiratory Coronavirus Syndrome (MERS-CoV) - are the main causes of SARS-CoV outbreaks of Acute Respiratory Syndrome in 2002 and 2003 in Guangzhou Province, China (5-7). The MERS-CoV pathogen was responsible for the outbreak of severe respiratory diseases in 2012 in the Middle East (8).

Due to the high prevalence and wide distribution of coronaviruses, the high genetic of and recombination their diversity genomes, and the increase in human-animal activity, new coronaviruses due to periods of cross-infections of various species and sudden events occur periodically in humans (9). In late December of 2019, several local health centers reported several patients with pneumonia who were epidemiologically related to the wholesale market of marine and wet animals in Wuhan, Hubei Province, China (4, 9). As of July 8, 2020, infection with (SARS-CoV-2) has resulted in more than 11 million laboratory-confirmed cases and 530,000 confirmed deaths (10).

The findings show that COVID-19 is mild in 81% of patients and improves at home. In 14% of cases, the person develops severe symptoms, including pneumonia and shortness of breath. In 5% of cases, the patient's condition worsens, which is associated with respiratory failure, infectious shock and failure in other organs of the body. In general, the mortality rate of this virus is 3.4% and the results show that the mortality rate of this virus in the elderly and people with underlying diseases is significantly higher than healthy people (11). Examining the level of knowledge and attitude of health workers, they concluded that their level of knowledge about the use of personal equipment, infection control and isolation measures is poor (12-18).

Poor understanding of the disease among

health care workers can lead to a rapid spread of infections in delayed diagnosis and treatment. More than 100 health workers have lost their lives to COVID-19, a catastrophe for the world and an obstacle to the fight against disease (19). Guidelines for health care workers and online refresher courses have been developed by the World Health Organization, the Centers for Disease Control and various government agencies in different countries to enhance knowledge and prevention strategies (20).

There is a lack of studies on the knowledge and attitudes of health care workers towards COVID-19. However, a study of health care workers, mostly Asian and medical students, found that they did not know enough about COVID-19 but had a positive attitude toward preventing COVID-19 transmission (21).

There is currently no vaccine available for Covid 19 in Iran and the only way to combat it is to prevent this disease. Due to the fact that the prevalence of this virus is very high among the community and the strong spreading potential of this virus, especially in health care centers and the death of several medical colleagues as well as the lack of a study on the attitude and knowledge of staff in Iranian nurses, the aim of the study was to compare the knowledge and attitudes of nurses in emergency departments and special wards of medical training centers in Mazandaran province about the new coronavirus in 2020.

## **Methods**

This descriptive analytical cross-sectional study was performed in multispecialty hospitals in Sari, Iran, from the 28th April 2020 to the 25th August 2020.

In this study, a researcher-made questionnaire related to knowledge and attitude was used for data collection. The questionnaire consisted of 3 parts that addressed people's knowledge and attitudes about the new coronavirus. The first part included demographic data including age, sex, ward, hospital, education, and participants' knowledge of the new coronavirus. The second section assessed nurses' knowledge by asking questions about cause, incubation period, symptoms, risk group, consequences, source of transmission, prevention and treatment. The validity of questionnaire content was assessed by two methods measuring the content validity index (CVI) and content validity ratio (CVR) and by consulting the expert group (CVI=0.91 and CVR<0.64). In order to validate the questionnaire, 20 participants were given questions as a pilot to determine the level of acceptance and clarity of questions. The reliability of the questionnaire based on Cronbach's alpha was 85%. The scoring system of questionnaire consisted of: 2 points for each correct answer and 1 point for wrong answer. A total of 12 points of the total knowledge scores were considered sufficient. Participants were divided into two groups according to their level of knowledge: sufficient knowledge (points  $\geq 12$ ), and insufficient knowledge (points <12). The third part of the questionnaire evaluated the nurses' attitudes about the new corona using a set of 8 yes/no questions. Participants were divided into two groups according to their level of attitude: sufficient attitude (points  $\geq 6$ ), and insufficient attitude (points < 6).

The questionnaire was completed online and self-reportedly. After the research plan was approved by the research ethics committee of Mazandaran University of Medical Sciences, prepared researchers an online the questionnaire in Google Form. After stating the objectives of the study, participants were asked to complete the questionnaire if satisfied. Then, the link of the questionnaire was sent to nursing departments of emergency departments and intensive care of selected hospitals affiliated to Mazandaran University of Medical Sciences. Sampling was census, which was easily accessible and non-random. Samples included all nurses in the emergency and intensive care units of Mazandaran University of Medical Sciences who were willing to participate in this study. Exclusion criteria included those questionnaires that were not completed by 10% of the main questions.

## **Statistical analysis**

Data were entered into SPSS v.16. Mean and standard deviation and frequency distribution table were used to describe the data. The normality of the data was measured using the Kolmogorov-Smirnov test. To analyze the data from independent t-test (to compare the average knowledge and attitude by gender and ward) and analysis of variance (to compare the average knowledge and attitude by education) and correlation coefficient (relationship between knowledge and attitude and also Age and history with knowledge and attitude) were used and p-value less than 0.05 was considered as significant.

## **Results**

Ninety-two nurses participated in this study. The mean age of the subjects was  $36.35 \pm 8.81$  years (range 19 to 60) and most of them were female. The demographic information of nurses is shown in *Table 1*. In terms of work experience, they had an average of  $12.10 \pm 8.33$  years of experience.

Table 1. Demographic information of samples

Variable		No	%
Condon	Male	15	16.3
Gender	Female	77	83.7
Word	Special	43	46.7
ward	Emergency	49	53.3
Educational degree	Bachelor	74	81.3
	Master	14	15.4
	PhD	3	3.3
	Cyberspace	46	51.2
Source of	Articles	21	23.3
information about	Television	21	23.3
COVID-19	Conferences	r	2.2
	and Seminars	2	2.2

**Table 2** shows the mean and standard deviation of knowledge and attitude scores based on gender, type of department and education degree of nurses. The mean score of nurses' knowledge was  $7.78 \pm 0.85$ , showing they generally had insufficient knowledge (less than average). The mean score of their attitudes was  $6.87 \pm 0.90$ , which is at a sufficient level (above average). The results of the analysis showed that the average knowledge of women was higher than men, but there was no statistically

significant difference. The mean attitude of men was higher than women, however this difference was not significant. In addition, the average knowledge of nurses working in the intensive care unit was higher than the emergency department, but there was no statistically significant difference. The mean attitude of nurses in the intensive care unit was higher than the emergency department but the difference was not significant. In terms of education level, PhDs had the highest scores of knowledge and attitude, but the mean scores were not significantly related to the level of education.

**Table 3** examined the relationship between knowledge and attitude scores as well as age and history variables with knowledge and attitude. Analytical results showed that there was no statistically significant relationship between any types of the variables.

Variable		Knowledge		Attitude	
		Mean±SD	P-value	Mean±SD	P-value
Gender	Male	$7.78\pm0.89$	0.94	$7.06\pm0.96$	0.99
	Female	$7/80 \pm 0.85$		$6.83\pm0.89$	
Ward	Special	$7.80\pm0.75$	0.95	$7.05\pm0.78$	0.37
	Emergency	$7.73\pm0.93$		$6.73 \pm 0.99$	
Educational degree	Bachelor	$7.79\pm0.82$		$6.93\pm0.90$	
	Master	$7.57 \pm 1.01$	0.61	$6.69\pm0.85$	0.61
	PhD	$8.00\pm1.00$		$7.00 \pm 1.00$	

## Table 2. Knowledge and attitude based on gender, type of department and education

# Table 3. Relationship between knowledge, attitudeand the two variables of age and work experience

		The correlation coefficient	P- value
Knowledge	Attitude	0.01	0.90
	Age	0.01	0.90
	Work experience	-012	0.29
Attitude	Age	0.3	0.81
	Work experience	0.3	0.82

## **Discussion**

The aim of this study was to compare the knowledge and attitudes of nurses in emergency departments and special wards of training centers in Mazandaran province about the coronavirus in 2020.

Findings of this study showed that the mean of nurses' knowledge was  $7.78 \pm 0.85$  (insufficient knowledge) and the mean of nurses' attitude was  $6.87 \pm 0.90$  (sufficient attitude). Findings of the study indicated insufficient knowledge of nurses that is not consistent with other studies (22-24). Nurses' knowledge about the prevalence of COVID-19 directly affects the outcomes, prevention and control of Covid-19 infection (25). Therefore, since the sampling of this study was performed at the beginning of the Covid-19 epidemic, sufficient training had not yet been provided to staff on a regular basis.

In this study, 51% of nurses obtained their information through cyberspace, about 2.2% of nurses through conferences, 23.3% through articles, and 23.3% through radio and television. In another study, most health care providers obtained their information through media such as radio and television (26, 27). However, it is consistent with a study by Abdul Ghadir that most nurses had acquired their knowledge by the Internet (24). Abdul Ghadir was quoted by the New York Times as saying that Internet traffic in the United States has increased dramatically in recent years due to people searching for information about Covid-19.

In this study, due to nurses' access to cyberspace, even during treatment shifts, is one of the main reasons for obtaining information from the Internet.

The attitude of nurses was  $6.87\pm0.90$ , which was a sufficient and positive attitude and is consistent with the results of other studies (27, 28). There was no statistically significant difference between knowledge and attitude between emergency and special ward nurses. Considering that a study has not compared nurses so far, this lack of discrepancy between nurses' knowledge and attitude may be due to the small sample size.

There is no significant relationship between education level, age and background with knowledge and attitude which is consistent with Nemati et al.'s study that the total score of knowledge was not affected by age, level of education and work experience (23).

Our research has some limitations. First, no standard tools for assessing nurses' knowledge and attitudes have already been approved in COVID-19. Therefore, WHO, CDC guidelines and COVID-19 reports have been used. Second, knowledge and attitude were examined in the teaching hospitals of Mazandaran University of Medical Sciences and the results of this study may not reflect the nurses in the whole country. However, due to the fact that the web-based survey was conducted and the response rate is low (30%), this generalization limits the survey.

## Conclusion

The level of nurses' knowledge was insufficient and the level of their attitude was sufficient. Therefore, it is recommended that health policy makers and nursing managers hold training classes and webinars to increase the level of awareness among nursing staff.

## Ethical standards statement

This project has been done with the approval and financial support of the Research Council of Mazandaran University of Medical Sciences (IR.MAZUMS.REC.1399.043).

## Acknowledgments

It is necessary to sincerely thank and appreciate the research council and student research committee of Mazandaran University of Medical Sciences and the participants of the research project.

## Conflicts of interest

The authors have stated that no conflict of interest occurs.

## Authors' contributions

All authors have intellectually committed to the study design and process. The final manuscript was revised and accepted by all authors.

# References

1. Lipsitch M, Swerdlow DL, Finelli L. Defining the epidemiology of Covid-19 studies needed. N Engl J Med.2020;382:1194–1196.

2. Rismanbaf A. Potential treatments for COVID-19; a narrative literature review. Arch Acad Emerg Med.2020;8(1):e29

3. Zangrillo A, Beretta L, Silvani P, Colombo S, Scandroglio AM, Dell'Acqua A, et al. Fast reshaping of intensive care unit facilities in a large metropolitan Downloaded from militarymedj.ir at 16:00 +0430 on Wednesday August 5th 2020 [ DOI: 10.30491/JMM.22.2.184 ] hospital in Milan, Italy: facing the COVID-19 pandemic emergency. Critical care and resuscitation: journal of the Australasian Academy of Critical Care Medicine. 2020.

4. Remuzzi A, Remuzzi G. COVID-19 and Italy: what next?. Lancet (London, England). 2020.

5. Saffarinia, M. The prediction of mental health based on the anxiety and the social cohesion that caused by Coronavirus. Social Psychology Research, 2020; 9(36): 129-141.

6. Centers for Disease Control and Prevention. Quarantine andisolation. 2017.https://www.cdc.gov/quarantine/index. html(accessed Jan 30, 2020)

7. Barbisch D, Koenig KL, Shih FY. Is there a case for quarantine? Perspectives from SARS to Ebola. Disaster Med Public Health Prep2015; 9: 547–53.

8. Miles SH. Kaci Hickox: public health and the politics of fear. 2014.http://www.bioethics.net/2014/11/kacihickox-public-health-and-thepolitics-of-fear/ (accessed Jan 31, 2020).

9. Twu SJ, Chen TJ, Chen CJ, Olsen SJ, Lee LT, Fisk T, et al. Control measures for severe acute respiratory syndrome (SARS) in Taiwan. Emerg Infect Dis. 2003;9:718–20

10. WHO. Coronavirus disease (COVID-

19): situation report—170. July 8, 2020. https://www.who.int/docs/default-source/ coronaviruse/situation-reports/20200708-

covid-19-sitrep-170.pdf?sfvrsn=bca86036\_2 (accessed July 10, 2020)

11. Tavakoli A, Vahdat K, Keshavarz M. Novel Coronavirus Disease 2019 (COVID-19): An Emerging Infectious Disease in the 21st Century. Iran South Med J. 2020; 22 (6) :432-450. URL:

http://ismj.bpums.ac.ir/article-1-1222fa.html

12. Goldberg, D.P. & Hillier, V.F. (1979) A scaled version of the General Health Questionnaire. Psychological Medicine, 9, 139-145.

13. Taghavi S. Evaluation of Validity and Reliability of General Health Questionnaire (GHQ). Journal of psychology 2002;;5(4):381-98(Persian).

14. Gammon J, Hunt J, Musselwhite C. The stigmatisation of source isolation: a literature review. J Res Nurs 2019;24:677-693.

15. Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. Lancet 2020;395:912-920.

16. Purssell E, Gould D, Chudleigh J. Impact of isolation on hospitalised patients who are infectious: systematic review with meta-analysis. BMJ Open 2020;10:e030371.

17. Sharma A, Pillai DR, Lu M, Doolan C, Leal J, Kim J, et al. Impact of isolation precautions on quality of life: a metaanalysis. J Hosp Infect 2020;105:35-42.

Xiao H, Zhang Y, Kong D, Li S, Yang 18. N. Social Capital and Sleep Quality in Individuals Who Self-Isolated for 14 Days During the Coronavirus Disease 2019(COVID-19) Outbreak in January 2020 in China. Medical Science Monitor. 2020;26. MedScape. In Memoriam: Healthcare 19. Workers Who Have Died of COVID-19. Available (2020).online at: www.medscape.com/viewarticle/927976 (accessed April 06, 2020).

20. World Health Organisation. Emerging Respiratory Viruses, Including COVID-19: Methods for Detection, Prevention, Response and Control. (2020). Available online at: www.openwho.org/courses/introduction-toncov (accessed March 18, 2020).

21. Bhagavathula AS, Aldhaleei WA, Rahmani J, Mahabadi MA, Bandari DK. Novel coronavirus (COVID-19) knowledge and perceptions: a survey on healthcare workers. medRxiv. [Preprint]. (2020). doi: 10.2196/19160

22. Nemati M , Ebrahimi B, Nemati F. Assessment of Iranian Nurses' Knowledge and Anxiety Toward COVID-19 During the Current Outbreak in Iran. Arch Clin Infect Dis. 2020 April; 15(COVID-19):e102848

23. Jawed F, Manazir S, Zehra A, Riaz R. The novel Coronavirus disease (COVID-19) pandemic: Knowledge, attitude, practice, and perceived stress among health care workers in Karachi, Pakistan. Med J Islam Repub Iran. 2020;34:132.

24. Nashwan AJ, Abujaber AA, Mohamed AS, Villar RC, Al-Jabry MM. Nurses' willingness to work with COVID-19 patients: The role of knowledge and attitude. Nurs Open. 2020;00:1–7.

25. Zhang M, Zhou M, Tang F, Wang Y, Nie H, Zhang L, You G. Knowledge, attitude, and practice regarding COVID-19 among healthcare workers in Henan, China. J Hosp Infect. 2020 Jun;105(2):183-187. doi: 10.1016/j.jhin.2020.04.012. Epub 2020 Apr 9. PMID: 32278701; PMCID: PMC7194961. Bhagavathula AS, Aldhaleei WA, 26. Rahmani J, Mahabadi MA, Bandari DK. Novel coronavirus (COVID-19) knowledge and perceptions: a survey on healthcare workers. MedRxiv 2020. https:// doi.org/10.1101/2020.03.09.20033381.

27. Saqlain M, Munir MM, Rehman SU, Gulzar A, Naz S, Ahmed Z, Tahir AH, Mashhood M. Knowledge, attitude, practice and perceived barriers among healthcare workers regarding COVID-19: a crosssectional survey from Pakistan. J Hosp Infect. 2020;105(3):419-423. doi: 10.1016/j.jhin.2020.05.007. Epub 2020 May 11. PMID: 32437822; PMCID: PMC7211584. 28. Malik UR, Atif N, Hashmi FK, et al. Knowledge, Attitude, and Practices of Healthcare Professionals on COVID-19 and Risk Assessment to Prevent the Epidemic Spread: A Multicenter Cross-Sectional Study from Punjab, Pakistan. Int J Environ Res Public Health. 2020;17(17):6395. Published 2020 Sep 2. doi:10.3390/ijerph17176395